

## Cover Sheet

**DATE:** February 14, 2012

**SUBJECT:** Recommendations for Influenza Anti-viral Use

**INSTRUCTIONS:**

**REDISTRIBUTE AT YOUR DISCRETION.** Share this information with relevant SMEs or contacts (internal and external) as you see fit. **Remove this cover sheet before redistributing and replace it with your own.**

**For LOCAL HEALTH DEPARTMENT reference only**

DPHHS Subject Matter Resource for more information regarding this HAN, contact:

**DPHHS CDCP  
Epidemiology Section  
1-406-444-0273**

**DPHHS Health Alert Hotline:  
1-800-701-5769**

**DPHHS HAN Website:  
[www.han.mt.gov](http://www.han.mt.gov)**

**Remove this cover sheet before redistributing and replace it with your own.**

**Please ensure that DPHHS is included on your HAN distribution list.**

**Categories of Health Alert Messages:**

**Health Alert:** conveys the highest level of importance; warrants immediate action or attention.

**Health Advisory:** provides important information for a specific incident or situation; may not require immediate action.

**Health Update:** provides updated information regarding an incident or situation; unlikely to require immediate action.

**Please call DPHHS to update contact information at 444-0919**

# Information Sheet

**Date:** February 14, 2012

**Subject:** Recommendations for Influenza Anti-viral Use

**Information:**

In association with the increase in cases of influenza nationally, CDC has sent out guidance for use of antivirals in association with influenza. Healthcare providers are encouraged to test those patients presenting with an influenza-like-illness (ILI) and treat following the attached CDC recommendations. Patients requiring hospitalization with a severe ILI should have more confirmatory influenza testing performed at the Montana Public Health Laboratory to determine the best course of antiviral treatment. Local health jurisdictions and the medical community should also continue to offer the influenza vaccine to those unvaccinated individuals.

# This is an official CDC HAN Info Service Message

Distributed via the HAN Info Service  
February 09, 2012, 15:00 EST  
CDCHAN-00324-ADV-N

## CDC Recommendations for Influenza Antiviral Medications Remain Unchanged

A recent review of randomized clinical trial data for the influenza neuraminidase inhibitor antiviral medications, and two related commentaries, raised questions about the value of antiviral medications for the prevention and treatment of influenza. On February 7, 2012, the Centers for Disease Control and Prevention (CDC) posted a "Have You Heard?" piece in response. The CDC response states "After careful consideration of all available evidence, CDC guidance on the use of antiviral medications remains unchanged." CDC continues to recommend the use of neuraminidase inhibitor antiviral drugs (oral oseltamivir and inhaled zanamivir) as an important adjunct in the prevention and treatment of influenza.

These recommendations emphasize **early antiviral treatment** as soon as possible for patients who are severely ill and for those who are at greatest risk for complications from influenza.

Patients who should receive early antiviral treatment include:

- Hospitalized patients with suspected or confirmed influenza;
- Persons with severe or progressive illness;
- Outpatients who are at high risk for influenza complications (for example, young children, people 65 and older, pregnant women, and persons with certain underlying chronic medical conditions) (For a full list of people at high risk of influenza complications, see: [http://www.cdc.gov/flu/about/disease/high\\_risk.htm](http://www.cdc.gov/flu/about/disease/high_risk.htm)); and
- Persons with uncomplicated influenza who are not in a high risk group and who present within 48 hours of illness onset. These persons can be treated with antiviral medications based upon clinical judgment, because reviews of RCTs and observational studies have found consistent clinical benefit of early oseltamivir treatment in reducing the risk of lower respiratory tract complications such as those requiring antibiotics.

CDC estimates that influenza virus infections in the United States result in an average of more than 200,000 related hospitalizations and 3,300–49,000 deaths each year, depending upon the severity of the influenza season. Annual influenza vaccination of all persons aged 6 months and older is recommended, and is the best way to prevent influenza. However, available evidence for seasonal influenza and 2009 pandemic H1N1 consistently indicates that antiviral treatment, when initiated as soon as possible, can have clinical and public health benefit in reducing severe outcomes of influenza. Therefore, neuraminidase inhibitor antiviral medications continue to be recommended for treatment of influenza.

The full CDC response is available at:  
[http://www.cdc.gov/media/haveyouheard/stories/Influenza\\_antiviral.html](http://www.cdc.gov/media/haveyouheard/stories/Influenza_antiviral.html).

For a summary of CDC antiviral guidance for the 2011-2012 influenza season see:  
<http://www.cdc.gov/flu/pdf/professionals/antivirals/clinician-antivirals-2011.pdf>

For the full document outlining ACIP and CDC recommendations and references supporting the statements in the "Have You Heard," CDC response, see "Antiviral Agents for the Treatment and Chemoprophylaxis of Influenza. Recommendations of the Advisory Committee on Immunization Practices (ACIP)" at:  
<http://www.cdc.gov/mmwr/pdf/rr/rr6001.pdf>

To read the review of randomized clinical trial data for the influenza neuraminidase inhibitor antiviral medications and related commentaries, see:

- The Cochrane Library's "Neuraminidase inhibitors for influenza: methods change, principles don't" at:  
<http://www.thecochranelibrary.com/details/editorial/1442857/Neuraminidase-inhibitors-for-influenza-methods-change-principles-dont.html>
- The British Medical Journal's "Rethinking credible evidence synthesis" at:  
<http://www.bmj.com/content/344/bmj.d7898>
- The British Medical Journal's "Questions remain over safety and effectiveness of oseltamivir" at: <http://www.bmj.com/content/344/bmj.e467>

*The Centers for Disease Control and Prevention (CDC) protects people's health and safety by preventing and controlling diseases and injuries; enhances health decisions by providing credible information on critical health issues; and promotes healthy living through strong partnerships with local, national and international organizations.*

**DEPARTMENT OF HEALTH AND HUMAN SERVICES**

---

**Categories of Health Alert messages:**

- Health Alert** conveys the highest level of importance; warrants immediate action or attention.  
**Health Advisory** provides important information for a specific incident or situation; may not require immediate action.  
**Health Update** provides updated information regarding an incident or situation; unlikely to require immediate action.  
**HAN InfoService** provides general public health information; unlikely to require immediate action.

##This message was distributed to state and local health officers, public information officers, epidemiologists and HAN coordinators, as well as clinician organizations.##

=====  
You have received this message based upon the information contained within our emergency notification data base. If you have a different or additional e-mail or fax address that you would like us to use, please contact your state-based Health Alert Network program at your state or local health department.  
=====