


Montana Adolescent Immunization Coverage Rates, Ages 13-17, 2015-2018



At 11-12 years of age: CDC Recommends


- **Meningococcal ACWY** (one dose)
- **HPV** (two doses)
- **Tetanus, diphtheria, and whooping cough (pertussis) (Tdap)** (one dose)
- **Influenza (flu)** (every year)

≥1 Tdap: ≥1 dose of tetanus, diphtheria, acellular, pertussis vaccine at or after age 10 years; ≥1 MenACWY: ≥1 dose of quadrivalent meningococcal conjugate vaccine; ≥1 HPV: ≥1 dose of human papillomavirus vaccine; HPV Up to date: includes those with ≥ 3 doses, and those with 2 doses when the first HPV vaccine dose was initiated at age <15 years. Influenza coverage is by season which spans two calendar years. The 2015 rate listed represents the 2015-2016 influenza season; 2016 is for 2016-2017; 2017 is for 2017-2018; and 2018 is for 2018-2019.


How Montana compares to the U.S. Rates

Tdap, Influenza, MenACWY among Adolescents


Tdap



Influenza*




MenACWY*




HPV- Females

1 Dose




Up-To-Date




HPV- Males

1 Dose



Up-To-Date



*Indicates a statistically significant difference between the U.S. and Montana rates